



Name: \_\_\_\_\_ Date of Birth: \_\_\_\_\_

Allergies: \_\_\_\_\_

Weight: \_\_\_\_\_ kg \_\_\_\_\_ lbs. Height: \_\_\_\_\_ in. BSA: \_\_\_\_\_ m<sup>2</sup>  Actual  Adjusted

Diagnosis:  B-Cell Lymphoma  Other: \_\_\_\_\_

Start Date: \_\_\_\_\_ Cycle #: \_\_\_\_\_

Premedications			
Dexamethasone (Decadron)	10 mg IV push over 3 to 5 minutes, 30 minutes prior to rituximab-pvvr	Day 1 _____	
Diphenhydramine (Benadryl)	25 mg IV push over 2-3 minutes, 30 minutes prior to rituximab-pvvr	Day 1 _____	
Acetaminophen	650 mg PO 30 minutes prior to rituximab-pvvr	Day 1 _____	
Meperidine (Demerol)	25 mg IV push over 4 minutes, ONCE PRN for rigors related to rituximab-pvvr	Day 1 _____	
Drug/ Administration	Dose/Unit	Final Dose	Date
Rituximab-pvvr (Ruxience)* IV on Day 1 in NSS - Final concentration 1 mg/mL - Initiate first dose at 50 mg/hr for 30 min - If tolerated, may increase rate by 50 mg/hr every 30 min to maximum rate of 400 mg/hr - If first cycle tolerated, may begin subsequent cycles at 100 mg/hr, increasing by 100 mg/hr every 30 min to maximum rate of 400 mg/hr  <input type="checkbox"/> Dose capped _____ <input type="checkbox"/> Dose Reduction _____ % Reason: _____	375 mg/m <sup>2</sup>	_____ mg	Day 1 _____

\* Hepatitis B & C testing recommended for all patients prior to first rituximab-pvvr dose (Hepatitis B Surface Antigen; Hepatitis B Core Antibody, Hepatitis C Antibody)

Additional Orders

- Check vital signs every 15 minutes x 1 hour, then with each increase in rituximab-pvvr infusion; or every 30 minutes.
- PICC line insertion  Access port  CMP, Routine  CBC with differential, Routine

Physician Signature: \_\_\_\_\_ Date: \_\_\_\_\_ Time: \_\_\_\_\_